

available at [www.sciencedirect.com](http://www.sciencedirect.com)[www.elsevier.com/locate/brainres](http://www.elsevier.com/locate/brainres)**BRAIN  
RESEARCH**

## Research Report

# Exposure to 1800 MHz radiofrequency radiation induces oxidative damage to mitochondrial DNA in primary cultured neurons

Shangcheng Xu, Min Zhong<sup>1</sup>, Lei Zhang, Zhou Zhou, Wei Zhang, Yuan Wang, Xubu Wang, Maoquan Li, Yang Chen, Chunhai Chen, Mindi He, Guangbin Zhang, Zhengping Yu\*

Department of Occupational Health, Third Military Medical University, No.30 Gaotanyan Street, Shapingba District, Chongqing 400038, People's Republic of China

## ARTICLE INFO

## Article history:

Accepted 24 October 2009

## Keywords:

Radiofrequency radiation

Oxidative stress

Mitochondrial DNA

Neuron

Melatonin

## ABSTRACT

Increasing evidence indicates that oxidative stress may be involved in the adverse effects of radiofrequency (RF) radiation on the brain. Because mitochondrial DNA (mtDNA) defects are closely associated with various nervous system diseases and mtDNA is highly susceptible to oxidative stress, the purpose of this study was to determine whether radiofrequency radiation can cause oxidative damage to mtDNA. In this study, we exposed primary cultured cortical neurons to pulsed RF electromagnetic fields at a frequency of 1800 MHz modulated by 217 Hz at an average special absorption rate (SAR) of 2 W/kg. At 24 h after exposure, we found that RF radiation induced a significant increase in the levels of 8-hydroxyguanine (8-OHdG), a common biomarker of DNA oxidative damage, in the mitochondria of neurons. Consistent with this finding, the copy number of mtDNA and the levels of mitochondrial RNA (mtRNA) transcripts showed an obvious reduction after RF exposure. Each of these mtDNA disturbances could be reversed by pretreatment with melatonin, which is known to be an efficient in the brain. Together, these results suggested that 1800 MHz RF radiation could cause oxidative damage to mtDNA in primary cultured neurons. Oxidative damage to mtDNA may account for the neurotoxicity of RF radiation in the brain.

© 2009 Published by Elsevier B.V.

## 1. Introduction

The world-wide and rapidly growing use of mobile phones (MP) has raised considerable concerns about their potentially hazardous effects on human health. Compared to the other organs, the brain is exposed to relatively high-specific absorption rates (SAR) due to the close proximity of the mobile

telephone device to the head. Exposure to radiofrequency (RF) radiation emitting from MP has been postulated to result in a variety of neurological effects, including headaches, changes in sleep patterns, modifications in neuronal electrical activity, increases in the permeability of the blood-brain barrier (BBB) and disturbances in neurotransmitter release (Hossmann and Hermann, 2003; Repacholi, 1998; Valentini et al., 2007). There is

\* Corresponding author. Fax: +86 23 68752290.

E-mail address: [xsosci2009@yahoo.cn](mailto:xsosci2009@yahoo.cn) (Z. Yu).

Abbreviations: MP, mobile phone; RF, radiofrequency; SAR, average special absorption; 8-OHdG, 8-hydroxyguanine; BBB, blood-brain barrier; mtDNA, mitochondrial DNA; mtRNA, mitochondrial RNA; ROS, reactive oxygen species; H<sub>2</sub>O<sub>2</sub>, hydrogen peroxide; DCFH-DA, 2',7'-dichlorofluorescein diacetate

<sup>1</sup> Contributed equally to this study.


growing evidence to suggest that RF radiation induces oxidative stress in the brain and in other tissues (Ilhan et al., 2004; Irmak et al., 2002; Koylu et al., 2006; Moustafa et al., 2001; Oktem et al., 2005; Ozguner et al., 2005a,b; Sokolovic et al., 2008; Yurekli et al., 2006). Oxidative stress is known to be essential for pathophysiological processes in the brain, and it plays critical roles in cell apoptosis, general and specific gene expression, DNA damage, cell proliferation, inflammation process and mitochondrial dysfunction (Hayashi et al., 2008; Lin and Beal, 2006). The involvement of oxidative stress in manifold processes is compatible with the diverse nature of the observed effects induced by RF radiation (Balik et al., 2005; Cleary et al., 1996; Diem et al., 2005; Koylu et al., 2006; Schwarz et al., 2008; Zhao et al., 2007a,b). Thus, oxidative stress may be involved in the adverse effects of RF radiation on the nervous system.

Mitochondria are the major sites of production of reactive oxygen species (ROS). ROS are toxic by-products of respiration, and excess ROS are known to cause oxidation of unsaturated fatty acid, proteins, and DNA. Mitochondrial DNA (mtDNA) is particularly susceptible to oxidative stress for a variety of reasons, including its lack of protective histone-like proteins, its insubstantial capacity for repair following damage, and its proximity to the respiratory chain in the mitochondrial inner membrane (Evans et al., 2004). MtDNA damage is reflected by the presence of mtDNA mutation, by a decline in mtDNA copy number, and by a reduced number of mitochondrial RNA (mtRNA) transcripts (Ide et al., 2001). Each of these types of mtDNA damage can amplify oxidative stress by encoding the deficient critical proteins for the respiratory chain, which exacerbates further production of ROS and aggravates the oxidative damage to mitochondrial function (Evans et al., 2004; Wallace, 1999). Indeed, oxidative damage to mtDNA contributes to mitochondrial dysfunction in various mitochondrial-related diseases including aging, neurodegenerative diseases and ischemia. The presence of mtDNA mutations is closely related to a variety of neurological symptoms, such as migraine, dementia, and ataxia (Lin and Beal, 2006; Taylor and Turnbull, 2005; Xu et al., 2009).

Therefore, there is the possibility that RF radiation could cause oxidative damage to mtDNA in nerve cells, and this may account for the adverse effects of RF radiation on the nervous system. In order to address this, we exposed primary cultured cortical neurons to 1800 MHz radiofrequency radiation at the SAR value of 2 W/kg. We observed a striking degree of oxidative damage to mtDNA in these neurons. In addition, we discovered that these oxidative injuries to mtDNA could be successfully prevented by melatonin, a pineal neurohormone with known antioxidant capacity especially in the brain (Lee et al., 2001; Sokolovic et al., 2008).

## 2. Results

In the RF radiation group, exposure to RF radiation for a period of 24 h significantly increased the production of ROS compared with that of the sham-control group ( $P < 0.01$ ; Fig. 1). A similar result was obtained with the hydrogen peroxide ( $H_2O_2$ ) group.  $H_2O_2$  was applied as a positive control for




**Fig. 1 – ROS production in primary cultured cortical neurons exposed to RF radiation. After exposure to RF radiation, neurons were incubated with DCFH-DA at 37 °C for 20 min. Fluorescence images were acquired with a Leica fluorescence microscope at 485 nm for excitation and 530 nm for emission. Fluorescence intensity for each dish was measured within 25×25 pixel regions of interest and analyzed for pixel intensity by NIH Image. Cellular fluorescence intensity was expressed as the multiple of the level in control groups. \*\* $p < 0.01$  versus sham-exposed control group and Melatonin group, # $p < 0.05$  versus RF radiation group. Values are means  $\pm$  SE,  $n = 6$ .**

measuring the effects of oxidative stress (Fig. 1). The increased amount of ROS in the radiation group could be reduced by pretreatment with melatonin ( $P < 0.05$ ; Fig. 1). These results indicated that RF radiation induced oxidative stress in neurons, and that this effect could be reversed by melatonin pretreatment.

To detect the oxidative damage of mtDNA that was induced by RF radiation, we performed immunostaining detection and HPLC analysis to determine the levels of 8-OHdG in mitochondria. 8-OHdG is the most prominent DNA damage, and it is often used as a biomarker for the oxidative damage of DNA. As shown in Fig. 2A, few 8-OHdG-immunopositive cells were observed in either the control or the melatonin group. However, after exposure to RF radiation, the number of 8-OHdG-positive cells significantly increased. We also observed that most of the 8-OHdG-positive cells were colocalized with MitoTracker Red CMXRos in the cytoplasm of the neurons (Fig. 2A). This result suggested that the RF radiation damaged DNA in the mitochondria of neurons. Consistent with the immunostaining detection results, HPLC analysis revealed that there was an approximately two-fold greater increase in the levels of 8-OHdG in mitochondria of the RF radiation group compared to those of the control and melatonin groups ( $P < 0.01$ ), and melatonin pretreatment almost completely attenuated this type of mtDNA oxidative damage (Fig. 2B).


It has been reported that mtDNA damage is not only reflected by an increase in mtDNA mutations, but also by a decline in mtDNA copy numbers and in reduction in the numbers of mtRNA (Ide et al., 2001). Our study showed that RF radiation reduced the mtDNA copy numbers by half (Fig. 3). In addition, the mtDNA transcript levels of ND1, COX I and ND6



**Fig. 2 – Increase of 8-OHdG levels in mitochondria after RF radiation. (A)** After exposure to RF radiation, neurons were incubated with MitoTracker Red CMXRos probe and then 8-OHdG monoclonal antibody. Confocal microscopic scanning photographs indicated the fluorescence signals of 8-OHdG immunostaining (blue) and MitoTracker Red CMXRos (red). The merged images of blue and red fluorescence appeared as a bluish red color. The presented image is the representative of three independent experiments. **(B)** For HPLC analysis, mtDNA in exposed neurons was extracted and digested to deoxynucleotides using nuclease P1 and ACP. The 8-OHdG content of mitochondria for each experimental group was measured by HPLC using an electrochemical detection system in experimental groups. 8-OHdG amount was expressed as the number of 8-OHdG molecules per 10<sup>6</sup> dG, determined simultaneously with a UV monitor coupled to the HPLC system. \*\**p* < 0.01 versus sham-exposed control group and Melatonin group, #*p* < 0.05 versus RF radiation group. Values are means ± SE, *n* = 3. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

145 in RF-irradiated cells were 49%, 28% and 64% lower than those  
 146 of the controls, respectively (Fig. 4). All of these reductions in  
 147 mtDNA copy number and mtDNA transcripts were signifi-

cantly prevented by melatonin pretreatment. These results 148  
 further proved that RF radiation damaged mtDNA in neurons 149  
 (Figs. 3 and 4). 150



**Fig. 3 – Decrease of mtDNA copy number after RF radiation.** Quantitative real-time PCR was used to measure mtDNA copy number. A specific fragment of Complex IV was designed for mtDNA copy number quantification. The amounts of mtDNA were normalized to the internal control,  $\beta$ -actin. \*\* $p < 0.01$  versus sham-exposed control group and melatonin group, # $p < 0.05$  versus RF radiation group. Values are means  $\pm$  SE,  $n = 6$ .


### 3. Discussion

Oxidative stress makes a substantial contribution to various pathological processes in the nervous system (Lin and Beal, 2006). As a result of its high metabolic rate, its inefficient oxidant defense mechanism and its diminished cellular turnover, the nervous system is particularly vulnerable to oxidative stress. Increasing evidence indicates that the radio-frequency fields of MP may affect brain biological activity by inducing oxidative stress (Ilhan et al., 2004; Koylu et al., 2006; Sokolovic et al., 2008; Zhao et al., 2007a). Consistent with this, we demonstrated that 1800 MHz RF radiation significantly damaged mtDNA integrity by increasing the production of ROS

in primary cultured cortical neurons. To our knowledge, this study is the first to demonstrate a relationship between RF radiation exposure and mtDNA damage in neurons.

Oxidative stress refers to an imbalance between the intracellular production of free radicals and cellular defense mechanisms (Tsutsui et al., 2006). In the present study, we showed that RF radiation obviously elevated the ROS production in cortical neurons. Consistent with previous studies, this indicated that RF radiation induced oxidative stress in neurons (Ilhan et al., 2004; Koylu et al., 2006; Sokolovic et al., 2008; Zhao et al., 2007a). An explanation for these findings could be that RF radiation produced an excess of ROS via inhibition of the mitochondrial respiratory chain, prolongation of the life-span of free radicals and impairment of the antioxidant defense system (Koylu et al., 2006; Moustafa et al., 2001). However, several studies have reported that radiation emitted from MP has no effect of inducing oxidative stress (Ferreira et al., 2006; Hook et al., 2004; Oktem et al., 2006; Simko et al., 2006; Zeni et al., 2007). For example, Hook et al. did not detect the induction of oxidative stress in J774.16 mouse macrophage cells that were exposed for 20–22 h at 0.8 W/kg to either frequency-modulated continuous-wave field at 835.62 MHz or code division multiple access field at 847.74 MHz. The discrepancies between our results and those of previous studies may be due to differences in the RF signals that were used, such as frequency, orientation, modulation, power density and duration of exposure (Ilhan et al., 2004; Simko, 2007; Zeni et al., 2007). Furthermore, even when the same experimental design is used, distinct cell types may react differently to the same stimulus (Schwarz et al., 2008; Simko, 2007).

MtDNA encodes 13 key subunits of the respiratory chain, and it is highly susceptible to oxidative damage. In our study, there was a significant increase in 8-OHdG in mitochondria after RF field exposure. Concomitant with this alteration, the mtDNA copy number and the mtRNA transcript levels also decreased after RF radiation. These reductions might result from a ROS-induced mutation at the origins of replication and transcription (Coskun et al., 2004; Ide et al., 2001). In addition, the expression of mitochondrial-encoded genes is largely



**Fig. 4 – Decrease of mtRNA transcripts levels after RF radiation.** Quantitative real-time PCR was used to measure the levels of mtRNA transcripts. The amounts of mtRNA were normalized to the internal control,  $\beta$ -actin. \*\* $p < 0.01$ , \* $p < 0.05$  versus sham-exposed control group and melatonin group, ## $p < 0.01$ , # $p < 0.05$  versus RF radiation group. Values are means  $\pm$  SE,  $n = 6$ .

regulated by the copy number of mtDNA (Kanki et al., 2004). Therefore, a decrease in the mtDNA copy number may result in a corresponding decrease in mtRNA transcripts, which could ultimately lead to mitochondrial dysfunction. Furthermore, oxidative stress could directly degrade mtDNA and mtRNAs by damaging the mtDNA nucleoid structure and the stability of mtRNAs (Demple, 2004; Kang et al., 2007; Rodriguez-Gabriel et al., 2003). Importantly, oxidative stress could be amplified by the damaged mtDNA through encoding of the deficient critical proteins for the respiratory chain (Evans et al., 2004). It is conceivable that RF radiation-induced oxidative stress caused mitochondrial dysfunction through damage to mtDNA. In turn, reduced mitochondrial function might further exacerbate the production of ROS, which could induce further oxidative damage to mtDNA. This vicious cycle might proceed until mitochondrial failure and apoptosis occurred. Indeed, recent data showed that exposure to radiofrequency radiation decreased cerebral cytochrome c oxidase activity and activated apoptosis signaling pathways in primary cultured neurons (Ammari et al., 2008; Zhao et al., 2007b).

Melatonin is a neurohormone that is synthesized in and released from the pineal gland during the dark period. It is capable of rapidly crossing the blood–brain barrier and accumulating in high concentrations in the cortical and hippocampal cells (Cabrera et al., 2000). Melatonin is considered to be a potent antioxidant that works to detoxify a variety of ROS and stimulate anti-oxidative enzymes in many pathophysiological states, particularly in the brain (Gupta et al., 2003). In our study, melatonin administration could effectively attenuate the oxidative damage of mtDNA. It further indicated that oxidative stress was the critical neurotoxic effect of RF radiation. This was consistent with the previous facts that melatonin improved oxidative phosphorylation and protected mtDNA against oxidative damage at the mitochondrial level by decreasing ROS production (Acuna-Castroviejo et al., 2001; Chen et al., 2005; Gurkok et al., 2009; Jou et al., 2007; Pappolla et al., 1999).

In conclusion, our study indicates that 1800 MHz RF radiation induced oxidative damage to mtDNA, but that melatonin prevented this damage. Based on the facts that mtDNA defects are the key factors in various mitochondria-associated behavioral symptoms and diseases of the nervous system (Dong et al., 2007; Lin and Beal, 2006; Loeb et al., 2005; Warner et al., 2004), oxidative damage to mtDNA may be an important mechanism underlying the adverse effects of RF radiation that is emitted from MP on brain activity. However, it may be difficult to directly extrapolate our results (which were obtained using rodent cells) to humans for different exposure conditions and SAR values. In future studies, to investigate the neurotoxicity of MP radiation in mtDNA defects, greater care will need to be taken to replicate the actual RF exposure conditions for humans.

## 4. Experimental procedures

### 4.1. Cell culture and RF radiation

Cortical neurons were dissected from newborn SD rats that were purchased from the animal center of Third Military

Medical University. All procedures were carried out in strict accordance with the guidelines of the Animal Care Committee. Minced neocortexes were digested with trypsin-EDTA at 37 °C for 20 min. Cells were plated onto 35 mm dishes that were pre-coated with poly-L-lysine-coated (Sigma, St. Louis, MO), at a density of  $5 \times 10^5$  cells/2 ml in DMEM/F-12 medium (Gibco BRL, Grand Island, NY) supplemented with 10% fetal bovine serum (HyClone, Logan, UT), 10% horse serum (HyClone, Logan, UT) and 50 mg/ml penicillin/streptomycin. After 24 h, cells were cultured in Neurobasal™ medium (Gibco BRL, Paisley, UK) containing  $1 \times B_{27}$  supplement (Invitrogen, Carlsbad, CA) and 0.5 mM L-glutamine.

On day 8, neurons were exposed to a carrier frequency of 1800 MHz intermittently (5 min fields on/10 min fields off) or sham-exposed for 24 h at an average SAR of 2 W/kg as previously described (Diem et al., 2005; Zhao et al., 2007a). The RF-exposure system used in these experiments is the sXc-1800 system (Zurich, Switzerland). This system is designed for the testing of RF electromagnetic field exposures from mobile communication devices in the 1800-MHz region such as the GSM digital communication system. The fields were amplitude-modulated by rectangular pulses with a repetition frequency of 217 Hz and a duty cycle of 1:8. The exposure system was composed of a RF generator, an arbitrary function generator, a narrow band amplifier and two rectangular waveguides. Exposed and sham-exposed cell dishes were simultaneously placed inside a commercial incubator in which the environmental conditions were constant (37 °C, 5% CO<sub>2</sub>, 95% humidity). During the exposure, the value of SAR was monitored and the temperature response of the medium was assessed as described previously (Sun et al., 2006). The temperature difference between sham- and RF-exposed cultures never exceeded 0.05 °C. To enable blind exposures, the entire setup was controlled with a computer, which automatically controlled the exposure parameters, including exposure strength (SAR) and exposure time.

The cultured cells were assigned to one of five groups: (1) sham-exposed control, (2) Melatonin, (3) RF radiation, (4) RF radiation+Melatonin (5) hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) exposure. In groups (2) and (4), melatonin (Sigma, St. Louis, MO) was dissolved in ethanol and then freshly diluted with cell medium before application. Melatonin was used 4 h prior to RF radiation at a concentration of 200 nM. In group (5), cells treated with 100 μM H<sub>2</sub>O<sub>2</sub> for 24 h were used as a positive control of oxidative stress. In groups (2) and (5), cells were put into the incubator for sham-exposure. Before the experiment, we also designed an incubator control and found there were no differences between the cells in the normal incubator and those in the sham-exposed waveguide at an average SAR of 2 W/kg for 24 h.

### 4.2. Assay of intracellular reactive oxygen species (ROS)

The determination of intracellular oxidant production was based on the oxidation of DCFH-DA (Beyotime Company, China). Briefly, exposed cells were incubated with DCFH-DA at 37 °C for 20 min. DCFH-DA passively diffused into the cells and was deacetylated by esterases to form nonfluorescent 2',7'-dichlorofluorescein (DCFH). In the presence of ROS,

317 DCFH reacted with ROS to form the fluorescent product DCF,  
318 which was trapped inside the cells. The amount of emitted  
319 fluorescence was correlated with the quantity of ROS in the  
320 cell. Fluorescence images were acquired using a Leica  
321 fluorescence microscope at 485 nm for excitation and  
322 530 nm for emission. The fluorescence intensity for each  
323 dish was measured within 25×25 pixel regions of interest  
324 and analyzed for pixel intensity by NIH Image. The  
325 experiment was repeated six times and cellular fluorescence  
326 intensity was expressed as the multiple of the level in  
327 control groups.

#### 328 4.3. Immunostaining of 8-OHdG in mitochondria

329 To measure the levels of 8-OHdG located in mitochondria,  
330 exposed cells were reacted with 200 nM MitoTracker Red  
331 CMXRos probe for 30 min at 37 °C according to the  
332 manufacturer's instructions. After being washed twice in  
333 cold PBS, cells were fixed with 2% (w/v) paraformaldehyde in  
334 PBS at 4 °C for 20 min. Then the cells were permeabilized  
335 with 0.5% Triton X-100 in PBS for 15 min at room  
336 temperature. The cells were washed twice with PBS, before  
337 being blocked with normal goat serum at 37 °C for 30 min.  
338 Subsequently, the cells were incubated with a mouse  
339 monoclonal antibody against 8-OHdG at 1:100 dilution  
340 overnight at 4 °C, and then with Cy5-labeled goat anti-  
341 mouse IgG (H+L) (Beyotime Company, China) at 1:200  
342 dilution for 1 h at 37 °C. The cells were mounted in mounting  
343 medium and visualized under a Leica confocal laser scanning  
344 microscope (TCS SP2, Germany). For two-color fluorescence,  
345 the red and blue signals were collected simultaneously and  
346 then superimposed. Colocalization appeared as a bluish red  
347 color due to mixing of the red and blue signals.

#### 348 4.4. HPLC analysis of 8-OHdG

349 To analyze the 8-OHdG contents in mitochondria, mtDNA  
350 was isolated from the neurons using a Qiagen Plasmid Midi  
351 kit, as previously described (Cahill et al., 1997). After  
352 mtDNA was digested with nuclease P1 and Escherichia  
353 coli alkaline phosphatase, the contents of 8-OHdG were  
354 assessed using high-performance liquid chromatography  
355 (HPLC) with an electrochemical detection system as  
356 reported. The 8-OHdG amount was expressed as the number  
357 of 8-OHdG molecules per 10<sup>6</sup> dG, determined simultaneously  
358 with a UV monitor coupled to the HPLC system (Cahill et al.,  
359 1999).

#### 360 4.5. Detection of mtDNA copy number and mtDNA 361 transcripts levels

362 We applied real-time PCR to detect mtDNA copy number and  
363 mtRNA transcripts levels as previously described (Xu et al.,  
364 2009). The real-time PCR was studied with the iQ<sup>TM</sup>5 Real-  
365 Time PCR Detection System (Bio-Rid, USA) and using the  
366 SYBR Green I detection method. Each real-time PCR reaction  
367 (25 µl total volume) contained 1 µl of template DNA (100 µg)  
368 or cDNA, 12.5 µl of 2×SYBR<sup>®</sup> Green Realtime PCR Master  
369 (Toyobo, Osaka, Japan), 1 µl of each of the forward and  
370 reverse primers and 9.5 µl ultrapure water. Samples were

denatured by heating at 95 °C for 4 min, followed by 40 cycles  
of amplification and quantification (95 °C for 15 s and 58 °C  
for 15 s) and by a final extension cycle (72 °C, 90 s). Each  
measurement was repeated six times and was normalized  
against the control.

For mtDNA copy number analysis, total cellular DNA from  
cells was extracted using the QIAamp DNA Mini kit (Qiagen,  
Hilden, Germany) according to the manufacturer's instruc-  
tions. As previously described, we compared relative amounts  
of mtDNA and nuclear DNA copy numbers. The mtDNA  
amplicons were generated from Complex IV segment. The  
nuclear amplicon was generated through amplification of a β-  
actin segment. The mtDNA primers (Accession No.  
NC\_001665) were 5-AGCAGGGA. TACCTCGTCGTTAC-3 and 5-  
GTTTTGATGCGAAGGCTTCTCAA-3 (fragment length,  
140 bp), and there were designed to minimize amplification  
of mtDNA pseudogenes embedded in the nuclear DNA. The  
primers for β-actin (accession no. NC\_005111), which were  
chosen as the internal standards, were 5-TCCCAGCACACTTA  
ACTTAGC-3 and 5-AGCCACAAGAACTCAGG-3 (fragment  
length, 98 bp). The threshold cycle number (Ct) values of β-  
actin and the mtDNA were determined for each individual  
quantitative PCR run. The -ddCt (mtDNA to β-actin) repre-  
sented the mtDNA copy number in a cell.

For mtRNA transcripts quantification, total RNA was  
extracted with TriPure (Roche, Indianapolis, IN) reagent.  
Reverse transcription was performed with ReverTra Ace  
reverse transcriptase (Toyobo, Osaka, Japan). The primers for  
NADH dehydrogenase subunits 1 (Accession No. NC\_001665  
(2740-3696)) were 5-CGAATACGCCGAGGAC C-3 and 5-  
TGCTAGGAAAATTGGCAG GGA-3 (fragment length, 298 bp).  
The primers for cytochrome c oxidase subunits I (accession no.  
NC\_001665 (5323-6867)) were 5-AGCAGGGA-  
TACCTCGTCGTTAC-3 and 5-GTTTTGATGCGAAGGC-  
TTCTCAA-3 (fragment length, 140 bp). The primers for  
NADH dehydrogenase subunits 6 (accession no. NC\_001665  
(13543-14061)) were 5-GATTGATTGTTAGTGATGTATTG-3  
and 5-CTCAGTAGCCATAGCAGTTG-3 (fragment length,  
137 bp). As mentioned above, β-actin was selected as the  
internal standard. The threshold cycle number (Ct) values of  
β-actin and mtDNA transcripts were determined. The -ddCt  
(mtDNA transcripts to β-actin) represented the mtDNA  
transcripts in a cell.

#### 414 4.6. Statistical analysis

415 All the experimental data were expressed as the means±SE.  
416 Each experiment was carried out at least three times.  
417 Statistical significances of differences between two groups  
418 were determined using Student's t-test. All P values were two-  
419 sided, and P<0.05 was considered statistically significant.

#### 420 Acknowledgments

421 We appreciate Dr. Mengxia Li, Dr. Wen Zeng, Dr. Yu Xu, Dr. Min  
422 Li for their hospitable and professional technical assistance.  
423 Grant support was from National Natural Science Foundation  
424 of China (No. 30470441).  
425

## 42 6 REFERENCES

- Acuna-Castroviejo, D., Martin, M., Macias, M., Escames, G., Leon, J., Khaldy, H., Reiter, R.J., 2001. Melatonin, mitochondria, and cellular bioenergetics. *J. Pineal Res.* 30, 65–74.
- Ammari, M., Lecomte, A., Sakly, M., Abdelmelek, H., de-Seze, R., 2008. Exposure to GSM 900 MHz electromagnetic fields affects cerebral cytochrome c oxidase activity. *Toxicology* 250, 70–74.
- Balik, H.H., Turgut-Balik, D., Balikci, K., Ozcan, I.C., 2005. Some ocular symptoms and sensations experienced by long term users of mobile phones. *Pathol. Biol. (Paris)* 53, 88–91.
- Cabrera, J., Reiter, R.J., Tan, D.X., Qi, W., Sainz, R.M., Mayo, J.C., Garcia, J.J., Kim, S.J., El-Sokkary, G., 2000. Melatonin reduces oxidative neurotoxicity due to quinolinic acid: *in vitro* and *in vivo* findings. *Neuropharmacology* 39, 507–514.
- Cahill, A., Wang, X., Hoek, J.B., 1997. Increased oxidative damage to mitochondrial DNA following chronic ethanol consumption. *Biochem. Biophys. Res. Commun.* 235, 286–290.
- Cahill, A., Stabley, G.J., Wang, X., Hoek, J.B., 1999. Chronic ethanol consumption causes alterations in the structural integrity of mitochondrial DNA in aged rats. *Hepatology* 30, 881–888.
- Chen, L.J., Gao, Y.Q., Li, X.J., Shen, D.H., Sun, F.Y., 2005. Melatonin protects against MPTP/MPP+ -induced mitochondrial DNA oxidative damage *in vivo* and *in vitro*. *J. Pineal Res.* 39, 34–42.
- Cleary, S.F., Du, Z., Cao, G., Liu, L.M., McCrady, C., 1996. Effect of isothermal radiofrequency radiation on cytolytic T lymphocytes. *FASEB J.* 10, 913–919.
- Coskun, P.E., Beal, M.F., Wallace, D.C., 2004. Alzheimer's brains harbor somatic mtDNA control-region mutations that suppress mitochondrial transcription and replication. *Proc. Natl. Acad. Sci. U. S. A.* 101, 10726–10731.
- Demple, B., 2004. Grasping the message: regulated mRNA stability in free radical stress responses. Rodriguez-Gabriel MA, Burns G, McDonald WH et al. RNA-binding protein Csx1 mediates global control of gene expression in response to oxidative stress. *EMBO J.* 2003 (22), 6256–6266 *Redox Rep.* 9, 3–5.
- Diem, E., Schwarz, C., Adlkofer, F., Jahn, O., Rudiger, H., 2005. Non-thermal DNA breakage by mobile-phone radiation (1800 MHz) in human fibroblasts and in transformed GFSH-R17 rat granulosa cells *in vitro*. *Mutat. Res., Genet. Toxicol. Environ. Mutagen.* 583, 178–183.
- Dong, W., Gao, D., Zhang, X., 2007. Mitochondria biogenesis induced by resveratrol against brain ischemic stroke. *Med. Hypotheses* 69, 700–701.
- Evans, M.D., Dizdaroglu, M., Cooke, M.S., 2004. Oxidative DNA damage and disease: induction, repair and significance. *Mutat. Res.* 567, 1–61.
- Ferreira, A.R., Bonatto, F., de Bittencourt Pasquali, M.A., Polydoro, M., Dal-Pizzol, F., Fernandez, C., de Salles, A.A., Moreira, J.C., 2006. Oxidative stress effects on the central nervous system of rats after acute exposure to ultra high frequency electromagnetic fields. *Bioelectromagnetics* 27, 487–493.
- Gupta, Y.K., Gupta, M., Kohli, K., 2003. Neuroprotective role of melatonin in oxidative stress vulnerable brain. *Indian J. Physiol. Pharmacol.* 47, 373–386.
- Gurkok, G., Coban, T., Suzen, S., 2009. Melatonin analogue new indole hydrazide/hydrazone derivatives with antioxidant behavior: synthesis and structure-activity relationships. *J. Enzyme Inhib. Med. Chem.* 24, 506–515.
- Hayashi, Y., Yoshida, M., Yamato, M., Ide, T., Wu, Z., Ochi-Shindou, M., Kanki, T., Kang, D.C., Sunagawa, K., Tsutsui, H., Nakanishi, H., 2008. Reverse of age-dependent memory impairment and mitochondrial DNA damage in microglia by an overexpression of human mitochondrial transcription factor A in mice. *J. Neurosci.* 28, 8624–8634.
- Hook, G.J., Spitz, D.R., Sim, J.E., Higashikubo, R., Baty, J.D., Moros, E. G., Roti Roti, J.L., 2004. Evaluation of parameters of oxidative stress after *in vitro* exposure to FMCW- and CDMA-modulated radiofrequency radiation fields. *Radiat. Res.* 162, 497–504.
- Hossmann, K.A., Hermann, D.M., 2003. Effects of electromagnetic radiation of mobile phones on the central nervous system. *Bioelectromagnetics* 24, 49–62.
- Ide, T., Tsutsui, H., Hayashidani, S., Kang, D., Suematsu, N., Nakamura, K., Utsumi, H., Hamasaki, N., Takeshita, A., 2001. Mitochondrial DNA damage and dysfunction associated with oxidative stress in failing hearts after myocardial infarction. *Circ. Res.* 88, 529–535.
- Ilhan, A., Gurel, A., Armutcu, F., Kamisli, S., Iraz, M., Akyol, O., Ozen, S., 2004. *Ginkgo biloba* prevents mobile phone-induced oxidative stress in rat brain. *Clin. Chim. Acta* 340, 153–162.
- Irmak, M.K., Fadillioglu, E., Gulec, M., Erdogan, H., Yagmurca, M., Akyol, O., 2002. Effects of electromagnetic radiation from a cellular telephone on the oxidant and antioxidant levels in rabbits. *Cell Biochem. Funct.* 20, 279–283.
- Jou, M.J., Peng, T.I., Yu, P.Z., Jou, S.B., Reiter, R.J., Chen, J.Y., Wu, H.Y., Chen, C.C., Hsu, L.F., 2007. Melatonin protects against common deletion of mitochondrial DNA-augmented mitochondrial oxidative stress and apoptosis. *J. Pineal Res.* 43, 389–403.
- Kang, D., Kim, S.H., Hamasaki, N., 2007. Mitochondrial transcription factor A (TFAM): roles in maintenance of mtDNA and cellular functions. *Mitochondrion* 7, 39–44.
- Kanki, T., Ohgaki, K., Gaspari, M., Gustafsson, C.M., Fukuoh, A., Sasaki, N., Hamasaki, N., Kang, D., 2004. Architectural role of mitochondrial transcription factor A in maintenance of human mitochondrial DNA. *Mol. Cell Biol.* 24, 9823–9834.
- Koylu, H., Mollaoglu, H., Ozguner, F., Naziroglu, M., Delibas, N., 2006. Melatonin modulates 900 MHz microwave-induced lipid peroxidation changes in rat brain. *Toxicol. Ind. Health* 22, 211–216.
- Lee, M.C., Chung, Y.T., Lee, J.H., Jung, J.J., Kim, H.S., Kim, S.U., 2001. Antioxidant effect of melatonin in human retinal neuron cultures. *Exp. Neurol.* 172, 407–415.
- Lin, M.T., Beal, M.F., 2006. Mitochondrial dysfunction and oxidative stress in neurodegenerative diseases. *Nature* 443, 787–795.
- Loeb, L.A., Wallace, D.C., Martin, G.M., 2005. The mitochondrial theory of aging and its relationship to reactive oxygen species damage and somatic mtDNA mutations. *Proc. Natl. Acad. Sci. U. S. A.* 102, 18769–18770.
- Moustafa, Y.M., Moustafa, R.M., Belacy, A., Abou-El-Ela, S.H., Ali, F.M., 2001. Effects of acute exposure to the radiofrequency fields of cellular phones on plasma lipid peroxide and antioxidant activities in human erythrocytes. *J. Pharm. Biomed. Anal.* 26, 605–608.
- Oktem, F., Ozguner, F., Mollaoglu, H., Koyu, A., Uz, E., 2005. Oxidative damage in the kidney induced by 900-MHz-emitted mobile phone: protection by melatonin. *Arch. Med. Res.* 36, 350–355.
- Oktem, G., Uslu, S., Vatanserver, S.H., Aktug, H., Yurtseven, M.E., Uysal, A., 2006. Evaluation of the relationship between inducible nitric oxide synthase (iNOS) activity and effects of melatonin in experimental osteoporosis in the rat. *Surg. Radiol. Anat.* 28, 157–162.
- Ozguner, F., Oktem, F., Armagan, A., Yilmaz, R., Koyu, A., Demirel, R., Vural, H., Uz, E., 2005a. Comparative analysis of the protective effects of melatonin and caffeic acid phenethyl ester (CAPE) on mobile phone-induced renal impairment in rat. *Mol. Cell Biochem.* 276, 31–37.
- Ozguner, F., Oktem, F., Ayata, A., Koyu, A., Yilmaz, H.R., 2005b. A novel antioxidant agent caffeic acid phenethyl ester prevents long-term mobile phone exposure-induced renal impairment in rat. Prognostic value of malondialdehyde, N-acetyl-beta-D-glucosaminidase and nitric oxide determination. *Mol. Cell Biochem.* 277, 73–80.

- 563 Pappolla, M.A., Chyan, Y.J., Poeggeler, B., Bozner, P., Ghiso, J.,  
564 LeDoux, S.P., Wilson, G.L., 1999. Alzheimer beta protein  
565 mediated oxidative damage of mitochondrial DNA: prevention  
566 by melatonin. *J. Pineal Res.* 27, 226–229.
- 567 Repacholi, M.H., 1998. Low-level exposure to radiofrequency  
568 electromagnetic fields: health effects and research needs.  
569 *Bioelectromagnetics* 19, 1–19.
- 570 Rodriguez-Gabriel, M.A., Burns, G., McDonald, W.H., Martin, V.,  
571 Yates III, J.R., Bahler, J., Russell, P., 2003. RNA-binding protein  
572 Csx1 mediates global control of gene expression in response to  
573 oxidative stress. *EMBO J.* 22, 6256–6266.
- 574 Schwarz, C., Kratochvil, E., Pilger, A., Kuster, N., Adlkofer, F.,  
575 Rudiger, H.W., 2008. Radiofrequency electromagnetic fields  
576 (UMTS, 1,950 MHz) induce genotoxic effects *in vitro* in human  
577 fibroblasts but not in lymphocytes. *Int. Arch. Occup. Environ.*  
578 *Health* 81, 755–767.
- 579 Simko, M., 2007. Cell type specific redox status is responsible for  
580 diverse electromagnetic field effects. *Curr. Med. Chem.* 14,  
581 1141–1152.
- 582 Simko, M., Hartwig, C., Lantow, M., Lupke, M., Mattsson, M.O.,  
583 Rahman, Q., Rollwitz, J., 2006. Hsp70 expression and free  
584 radical release after exposure to non-thermal radio-frequency  
585 electromagnetic fields and ultrafine particles in human Mono  
586 Mac 6 cells. *Toxicol. Lett.* 161, 73–82.
- 587 Sokolovic, D., Djindjic, B., Nikolic, J., Bjelakovic, G., Pavlovic, D.,  
588 Kocic, G., Krstic, D., Cvetkovic, T., Pavlovic, V., 2008. Melatonin  
589 reduces oxidative stress induced by chronic exposure of  
590 microwave radiation from mobile phones in rat brain. *J. Radiat.*  
591 *Res. (Tokyo)* 49, 579–586.
- 592 Sun, L.X., Yao, K., Wang, K.J., Lu, D.Q., Hu, H.J., Gao, X.W.,  
593 Wang, B.H., Zheng, W., Lou, J.L., Wu, W., 2006. Effects of  
594 1.8 GHz radiofrequency field on DNA damage and expression  
595 of heat shock protein 70 in human lens epithelial cells.  
596 *Mutat. Res., Fundam. Mol. Mech. Mutagen.* 602,  
597 135–142.
- Taylor, R.W., Turnbull, D.M., 2005. Mitochondrial DNA mutations 598  
in human disease. *Nat. Rev., Genet.* 6, 389–402. 599
- Tsutsui, H., Ide, T., Kinugawa, S., 2006. Mitochondrial oxidative 600  
stress, DNA damage, and heart failure. *Antioxid. Redox Signal.* 601  
8, 1737–1744. 602
- Valentini, E., Curcio, G., Moroni, F., Ferrara, M., De Gennaro, L.,  
603 Bertini, M., 2007. Neurophysiological effects of mobile phone  
604 electromagnetic fields on humans: a comprehensive review.  
605 *Bioelectromagnetics* 28, 415–432. 606
- Wallace, D.C., 1999. Mitochondrial diseases in man and mouse. 607  
*Science* 283, 1482–1488. 608
- Warner, D.S., Sheng, H., Batinic-Haberle, I., 2004. Oxidants,  
609 antioxidants and the ischemic brain. *J. Exp. Biol.* 207, 3221–3231. 610
- Xu, S.C., Zhong, M., Zhang, L., Wang, Y., Zhou, Z., Hao, Y.T., Zhang,  
611 W.Y., Yang, X.S., Wei, A.M., Pei, L.P., Yu, Z.P., 2009.  
612 Overexpression of Tfam protects mitochondria against  
613 beta-amyloid-induced oxidative damage in SH-SY5Y cells.  
614 *FEBS J.* 276, 4224–4233. 615
- Yurekli, A.I., Ozkan, M., Kalkan, T., Saybasili, H., Tuncel, H.,  
616 Atukeren, P., Gumustas, K., Seker, S., 2006. GSM base station  
617 electromagnetic radiation and oxidative stress in rats.  
618 *Electromagn. Biol. Med.* 25, 177–188. 619
- Zeni, O., Di Pietro, R., d'Ambrosio, G., Massa, R., Capri, M.,  
620 Naarala, J., Juutilainen, J., Scarfi, M.R., 2007. Formation of  
621 reactive oxygen species in L929 cells after exposure  
622 to 900 MHz RF radiation with and without co-exposure to  
623 3-chloro-4-(dichloromethyl)-5-hydroxy-2(5H)-furanone.  
624 *Radiat. Res.* 167, 306–311. 625
- Zhao, R., Zhang, S., Xu, Z., Ju, L., Lu, D., Yao, G., 2007a. Studying  
626 gene expression profile of rat neuron exposed to 1800 MHz  
627 radiofrequency electromagnetic fields with cDNA microassay.  
628 *Toxicology* 235, 167–175. 629
- Zhao, T.Y., Zou, S.P., Knapp, P.E., 2007b. Exposure to cell phone  
630 radiation up-regulates apoptosis genes in primary cultures of  
631 neurons and astrocytes. *Neurosci. Lett.* 412, 34–38. 632